



**United Nations
Environment Programme**

**Food and Agriculture Organization
of the United Nations**

Distr.: General
1 December 2005

English only

**Rotterdam Convention on the Prior Informed
Consent Procedure for Certain Hazardous
Chemicals and Pesticides in International Trade
Chemical Review Committee**

Second meeting

Geneva, 13–17 February 2006

Item 5 (b) of the provisional agenda*

**Inclusion of chemicals in Annex III of the Rotterdam Convention:
review of notifications of final regulatory actions to ban
or severely restrict a chemical: Endosulfan**

Endosulfan: supporting documentation provided by Thailand

Note by the secretariat

The annex to the present note contain the supporting documentation provided by Thailand in support of its final regulatory action on endosulfan.


* UNEP/FAO/RC/CRC.2/1.

Annex

List of supporting documentation on Endosulfan submitted by Thailand

- Pesticide Manual 13th edition
-

IRAC 2A; cyclodiene



whitefly parasite

r.h. The product

his crop black
asshouse whitefly
d as the parasitic
g the 1930s and
whitefly infestations
mosa is one of the
rers Applied
t; M&R Durango;
e.

and feeding upon
e, laying its egg in
ed up to a range of
axi
2. Dispersal flights
is introduced into
tised pupae in
for attachment to
ass successfully,
ts of tomatoes,
exceptions,
d when the pest is
t pupal parasites
types Parasitised
of other
fungicides.

eye or skin
have been

NOMENCLATURE: Common name endosulfan (BSO, E-ISO, (m) F-ISO, ANSI, ESA); thiodan (Iran, USSR); benzoepin (JMAF); no name (Italy)

IUPAC name (1,4,5,6,7,7-hexachloro-8,9,10-trinorborn-5-en-2,3-ylenebismethylene) sulfite;
6,7,8,9,10,10-hexachloro-1,5,5a,6,9,9a-hexahydro-6,9-methano-2,4,3-benzodioxathiepine 3-oxide
Chemical Abstracts name 6,7,8,9,10,10-hexachloro-1,5,5a,6,9,9a-hexahydro-6,9-methano-2,4,3-benzodioxathiepine 3-oxide

CAS RN [115-29-7] endosulfan; [959-98-8], formerly [33213-66-0] alpha-endosulfan; [33213-65-9], formerly [891-86-1] and [19670-15-6] beta-endosulfan **EC no.** 204-079-4
Development codes Hoe 02 671 (Hoechst); FMC 5462 **Official codes** OMS 204 (α); OMS 205 (β); OMS 570; ENT 23979

PHYSICAL CHEMISTRY: **Composition** Endosulfan is a mixture of two stereoisomers: alpha-endosulfan, endosulfan (I), stereochemistry 3α,5aβ,6α,9α,9aβ-, comprises 64–67% of the tech. grade; beta-endosulfan, endosulfan (II), stereochemistry 3α,5aα,6β,9β,9aα-, comprises 29–32%. Earlier reports on the stereochemistry of these isomers gave conflicting accounts (W. Riemschneider, *World Rev. Pest Control*, 1963, 2(4), 29). **Mol. wt.** 406.9 **M.f.** C₉H₆Cl₆O₃S **Form** Colourless crystals; (tech., cream to brown, mostly beige). **M.p.** ≥80°C (tech.); α- 109.2°C; β- 213.3°C **V.p.** 0.83 mPa (20°C) for 2:1 mixture of α- and β- isomers **K_{ow} logP** for α- = 4.74; β- = 4.79 (both at pH 5) **Henry** α- 1.48; β- 0.07 (both Pa m³ mol⁻¹, 22°C, calc.) **S.g./density** c. 1.8 (20°C) (tech.) **Solubility** In water alpha-endosulfan 0.32, beta-endosulfan 0.33 (both in mg/l, 22°C). In ethyl acetate, dichloromethane, toluene 200, ethanol c. 65, hexane c. 24 (all in g/l, 20°C). **Stability** Stable to sunlight. Slowly hydrolysed in aqueous acids and alkalis, with the formation of the diol and sulfur dioxide.

COMMERCIALISATION: **History** Insecticide reported by W. Finkenbrink (*Nachrichtenbl. Dtsch. Pflanzenschutzdienstes (Braunschweig)*, 1956, 8, 183). Introduced by Hoechst AG (now Bayer CropScience) and, in the USA, by FMC Corp. **Patents** DE 1015797; US 2799685; GB 810602 (all to Hoechst) **Manufacturers** Aako; Bayer CropScience; Drexel; Excel; Hindustan; Makhteshim-Agan; Milenia; Parry; Seo Han; Sharda.

APPLICATIONS: **Biochemistry** Antagonist of the GABA receptor-chloride channel complex. **Mode of action** Non-systemic insecticide and acaricide with contact and stomach action. **Uses** Control of sucking, chewing, and boring insects and mites on a very wide range of crops, including fruit (including citrus), vines (0.1–0.2%), olives, vegetables including potatoes (0.1–0.2%), ornamentals, cucurbits, cotton (1.5–2.5 l/ha), tea (1.0–2.0 l/ha), coffee (1.5–2.0 l/ha), rice (1.5–2.0 l/ha), cereals (1.0–2.0 l/ha), maize and sorghum (1.5–2.5 l/ha), oilseed crops (1.0–2.0 l/ha), hops, hazels and sugar cane (1.5–2.5 l/ha), tobacco (0.8–1.5 l/ha), alfalfa, mushrooms, forestry, glasshouse crops, etc. Also controls tsetse flies. **Phytotoxicity** Glasshouse geraniums and chrysanthemums, alfalfa, and lima beans may be injured. **Formulation types** DP; EC; GR; SC; UL; WP; Powder concentrate. **Compatibility** Incompatible with strongly alkaline

endosulfan 363

pesticide manual 13th EDITION

Main Entries

materials. Selected products 'Fan' (FMC); 'Afidanil' (Efthymiadis); 'Algodán 350' (Ingeniería Industrial); 'Cekulfan' (Cequisa); 'Davonil' (Vapco); 'Devisulfan' (Devidayal); 'Endocel' (Excel); 'Endodhan' (Dhanuka); 'Endosol' (Aimco); 'Endostar' (Shaw Wallace); 'Hildan' (Hindustan); 'Lucasulfan' (Lucava); 'Mentor' (Crop Health); 'Phaser' (Bayer CropScience); 'Speed' (Nagarjuna Agrichem); 'Thiodan' (Bayer CropScience); 'Thionex' (Makhteshim-Agan); mixtures 'Tomahawk' (+ ethion) (Calliope).

ANALYSIS: Product analysis by glc (CIPAC Handbook, 1985, 1C, 2110; AOAC Methods, 17th Ed., 983.08). Residues determined by glc with MCD (*ibid.*, 976.23; *Pestic. Anal. Man.*, 1979, 201-A, 201-G, 201-I; 405; A. Ambrus et al., *J. Assoc. Off. Anal. Chem.*, 1981, 64, 773; *Man. Pestic. Residue Anal.*, 1987, 1, 5, 6, S19; *Anal. Methods Residues Pestic.*, 1988, Part I, M1, M12). Endosulfan isomers in drinking water by glc with ECD (AOAC Methods, 17th Ed., 990.06). Further methods available on request from Bayer CropScience.

MAMMALIAN TOXICOLOGY: Reviews FAO/WHO 83, 85 (see part 2 of the Bibliography). Oral Acute oral LD₅₀ for rats 70 mg (in aqueous suspension)/kg, 110 mg tech. (in oil)/kg, 76 mg alpha- isomer/kg, 240 g beta- isomer/kg; for dogs 77 mg tech./kg. Skin and eye Acute percutaneous LD₅₀ for rabbits 359 mg (in oil)/kg; for male rats >4000, female rats 500 mg/kg. Inhalation LC₅₀ (4 h) for male rats 0.0345, female rats 0.0126 mg/l. NOEL (2 y) for rats 15 ppm diet; (1 y) for dogs 10 ppm diet. ADI (JMPR) 0.006 mg/kg b.w. [1998]. Toxicity class WHO (a.i.) II; EPA (formulation) I (tech.) EC classification T; R24/25| Xi; R36| N; R50, R53

ECOTOXICOLOGY: Birds Acute oral LD₅₀ for mallard ducks 205–245, ring-necked pheasants 620–1000 mg/kg. Fish Highly toxic (LC₅₀ (96 h) for golden orfe 0.002 mg/l water) but, in practical use, should be harmless to wildlife. Daphnia LC₅₀ (48 h) 75–750 µg/l. Algae EC₅₀ (72 h) for green algae >0.56 mg/l. Bees Not toxic to bees under field conditions at an application rate of 1.6 l/ha (560 g endosulfan/ha). Worms NOEC 0.1 mg/kg dry weight.

ENVIRONMENTAL FATE: EHC 40 (WHO, 1984). Animals The principal route of elimination is faeces; most of the radioactivity is excreted within the first 48 hours. The amounts excreted are independent of dose level, number of dosages and isomerism. There are indications of species-specificity. Residues of endosulfan accumulate in the kidneys rather than in fat. Elimination from the kidneys takes place with DT₅₀ 7 d, but there is no sign of accumulation in the kidneys even after long-term feeding. Endosulfan is metabolised rapidly in mammalian organisms to less-toxic metabolites and to polar conjugates. Plants The plant metabolites (mainly endosulfan sulfate) were also found in animals and have thus been investigated from a toxicological point of view. 50% of residues are lost in 3–7 days (depending on plant species). Soil/Environment Endosulfan (alpha- and beta-) is degraded in soil with DT₅₀ 30–70 d. The main metabolite usually found was endosulfan sulfate, which is degraded more slowly and is, for this reason, the most important metabolite. DT₅₀ for total endosulfan (alpha- and beta- endosulfan and endosulfan sulfate) in the field is 5–8 mo. No leaching tendency was observed. K_{OC} 3000–20 000; K_d <3%.